

**School:**  
**Emergency Plan- Anaphylaxis**  
**School Year – 2024-25**

Student:  
 Contact:

Birthdate:

Grade/Teacher:

Student Specific Emergency/Diagnosis: **Anaphylaxis**  
 \*Epi should be sent on all field trips.

If you see this	Do this
<b>EPI KEPT IN OFFICE</b>	
1. Known or suspected exposure to allergen to: 2. Symptoms such as: <ul style="list-style-type: none"> <li>• LIST STUDENT SPECIFIC REACTIONS SUCH AS:               <ul style="list-style-type: none"> <li>*Mouth: itching and swelling of lips, tongue or mouth</li> <li>*Throat: itching and or tightness in throat, hoarseness</li> <li>*Skin: hives, itchy rash; swelling of face, around eyes, or extremities</li> <li>*Gut: nausea, abdominal cramps, vomiting, and/or diarrhea</li> <li>*Lung: shortness of breath, repetitive coughing or wheezing, nasal discharge</li> <li>*Heart: thread, pulse, passing out</li> </ul> </li> </ul>	1. Administer epi pen immediately as directed, see attached. Epi pen in office. <ul style="list-style-type: none"> <li>• To administer:</li> <li>• Position child to access thigh area</li> <li>• Pull off blue safety cap</li> <li>• Place orange tip of EpiPen on the thigh, at a right angle to the leg.</li> <li>• Swing and firmly push orange tip against outer thigh so it “clicks” and hold for a count of 10.</li> <li>• Remove EpiPen and discard in sharps container.</li> <li>• Massage injection site for 10 seconds.</li> </ul> 2. Call 911 3. Call eCare 4. Remain with student 5. Monitor respiratory status and vitals per nurse direction 6. Be prepared to administer second dose as directed 7. Allow a position of comfort
1. If unresponsive and not breathing	2. Begin CPR 3. Call 911 4. Get an AED 5. Call eCare

**COMMENTS:**

**Created:**